



A NEAR CYCLIC (m_1, m_2, \dots, m_r) -CYCLE SYSTEM OF COMPLETE MULTIGRAPH

Mowafaq Alqadri and Haslinda Ibrahim

School of Quantitative Sciences

College of Arts and Sciences

Universiti Utara Malaysia

06010 Sintok, Kedah, Malaysia

e-mail: moufaqq@yahoo.com

linda@uum.edu.my

Abstract

Let v, λ be positive integers, λK_v denote a complete multigraph on v vertices in which each pair of distinct vertices joining with λ edges. In this article, difference method is used to introduce a new design that decomposes $4K_v$ into cycles, when $v \equiv 2, 10 \pmod{12}$. This design merging between cyclic (m_1, \dots, m_r) -cycle system and near-four-factor is called a near cyclic (m_1, \dots, m_r) -cycle system.

1. Introduction

In this paper, it is considered that all graphs are undirected with no loops and vertices set Z_v . We denote the complete graph on v vertices by K_v . An m -cycle (respectively, m -path), denoted by (c_0, \dots, c_{m-1}) (respectively, $[c_0, \dots, c_{m-1}]$), consists of m distinct vertices $\{c_0, c_1, \dots, c_{m-1}\}$ and m edges

Received: October 22, 2016; Revised: January 27, 2017; Accepted: February 6, 2017

2010 Mathematics Subject Classification: 05C38, 05C70.

Keywords and phrases: (cyclic) m -cycle system, difference set, near-four-factor, parallel cycles.

$\{c_i c_{i+1}\}$, $0 \leq i \leq m-2$ and $c_0 c_{m-1}$ (respectively, $m-1$ edges $\{c_i c_{i+1}\}$, $0 \leq i \leq m-2$).

An (m_1, \dots, m_r) -cycle is the union of all edges in each m_i -cycle, $1 \leq i \leq r$. A decomposition of a graph G is a set of subgraphs $\{H_1, \dots, H_r\}$ of G whose edges set partitions the edge set of G . If K_v has a decomposition into r cycles of length m_1, m_2, \dots, m_r , then it is said an (m_1, \dots, m_r) -cycle system of order v that is defined as a pair (V, C) such that $V = V(K_v)$, and C is a collection of edge-disjoint m_i -cycles, for $1 \leq i \leq r$, which partitions the $E(K_v)$. In particular, if $m_1 = \dots = m_r = m$, then it is called an m -cycle system of order v or (K_v, C_m) -design.

A complete multigraph of order v , denoted by λK_v , can be obtained by replacing each edge of K_v with λ edges. A (m_1, \dots, m_r) -cycle system of λK_v is a pair (V, C) , where $V = V(\lambda K_v)$ and C is a collection of edge-disjoint m_i -cycles for $1 \leq i \leq r$ which partitions the edge multiset of λK_v . An automorphism of (m_1, \dots, m_r) -cycle system of λK_v is a bijection $\alpha : V(Z_v) \rightarrow V(Z_v)$ such that for any $(c_0, \dots, c_{t-1}) \in C$ if and only if $(\alpha(c_0), \dots, \alpha(c_{t-1})) \in C$, (m_1, \dots, m_r) -cycle system of λK_v is called *cyclic* if it has automorphism that is a permutation consisting of a single cycle of order v , for instance, $\alpha = (0, 1, \dots, v-1)$ and is said to be *simple* if all its cycles are distinct.

Given an m -cycle $C_m = (c_0, c_1, \dots, c_{m-1})$, by $C_m + i$ we mean $(c_0 + i, c_1 + i, \dots, c_{m-1} + i)$, where $i \in Z_v$. Analogously, if $C = \{C_{m_1}, C_{m_2}, \dots, C_{m_r}\}$ is an (m_1, \dots, m_r) -cycle, then we use $C + i$ instead of $\{C_{m_1} + i, C_{m_2} + i, \dots, C_{m_r} + i\}$. A set of cycles that generates the cyclic (m_1, \dots, m_r) -cycle system of λK_v by repeated addition of 1 modular v which is called a *starter set* (briefly δ).

The study of (m_1, \dots, m_r) -cycle system of λK_v has been considered the

most important problems in graph decomposition. The important is case $\lambda = 1, m_1 = \dots = m_r = m$. The existence question for a (K_v, C_m) -design has been solved by Alspach and Gavlas [2] in the case of m odd and by Šajna [11] for m even. While the existence question for a cyclic m -cycle has been settled when $m = 3$ [8], 5 and 7 [10]. For m even and $v \equiv 1 \pmod{2m}$, a cyclic m -cycle system of order v was proved for $m \equiv 0, 2 \pmod{4}$ in [6, 9]. Recently, Bryant et al. [3] showed the necessary and sufficient conditions for decomposing K_v into r cycles of lengths m_1, m_2, \dots, m_r or into r cycles of lengths m_1, m_2, \dots, m_r and perfect matching. Thus, the Alspach's problem has been settled which was posed in 1981 [1]. More recently, it has been extended to this decomposition for the complete multigraph λK_v in [4].

A k -factor of a graph G is a spanning subgraph whose vertices have a degree k . While a near- k -factor is a subgraph in which all vertices have a degree k with exception of one vertex (isolated vertex) which has a degree zero.

Moreover, in [7], Matarneh and Ibrahim introduced the decomposition of a complete multigraph $2K_v$, when $v \equiv 0 \pmod{12}$, by combination of cyclic (m_1, m_2, \dots, m_r) -cycle system and near-two-factor. In our paper, we propose a new design for decomposing a complete multigraph $4K_v$ when $v \equiv 2, 10 \pmod{12}$. This is obtained by merging a cyclic (m_1, \dots, m_r) -cycle system and near-four-factors that is called a *near cyclic (m_1, \dots, m_r) -cycle system* denoted by $NCCS(4K_v, \delta)$. Thus, we present $NCCS(4K_v, \delta)$ as a $(v \times \lfloor \delta \rfloor)$ array satisfying the following conditions:

- the cycles in row r and column i form a near-four-factor with focus r ,
- the cycles associated with rows contain no repetitions.

The main result of this paper is the following:

Theorem 1.1. *There exists a full simple cyclic (m_1, \dots, m_r) -cycle system of $4K_v$, $NCCS(4K_v, \delta)$, when $v \equiv 2, 10 \pmod{12}$.*

2. Preliminaries

Throughout this paper, we use difference set method that will be clarified in this section to obtain the main results.

Let $G = K_v$, for $a, b \in V(K_v)$ and $a \neq b$, the difference d of pair $\{a, b\}$ is $|a - b|$ or $v - |a - b|$, whichever is smaller. We define the difference d of any edge $ab \in E(K_v)$ as $\min\{|a - b|, v - |a - b|\}$. So, the difference of any edge in $E(K_v)$ is not exceeding $\frac{v}{2}$, ($1 \leq d \leq \lfloor v/2 \rfloor$). Let $C_n = (a_0, a_1, \dots, a_{n-1})$ (respectively, $P_n = [a_0, a_1, \dots, a_{n-1}]$) be an n -cycle (respectively, n -path) of K_v , the list of differences from C_n is a multiset $D(C_n) = \{\min\{|a_i - a_{i-1}|, v - |a_i - a_{i-1}|\} | i = 1, 2, \dots, n\}$, where $a_0 = a_n$ (respectively, $D(P_n) = \{\min\{|a_i - a_{i-1}|, v - |a_i - a_{i-1}|\} | i = 1, 2, \dots, n-1\}$). The list difference from $\delta = \{C_{m_1}, \dots, C_{m_t}\}$ is the multiset $D(\delta) = \bigcup_{i=1}^t D(C_{m_i})$.

Definition 2.1. Given a complete multigraph λK_v , when v even. A set $\delta = \{C_{m_1}, \dots, C_{m_t}\}$ of cycles of λK_v is $(\lambda K_v, \delta)$ -difference system if $D(\delta) = \bigcup_{i=1}^t D(C_i)$ covers each element of $Z_v^* = Z_v - \{0\}$ exactly λ times and the middle difference $\left(\frac{v}{2}\right)$ appears $\left\{\frac{\lambda}{2}\right\}$ times.

As a particular result of the theory developed in [5], we have:

Proposition 2.1. A set $\delta = \{C_1, \dots, C_t\}$ of m_i -cycles, where $i = 1, 2, \dots, t$ is a starter set of a cyclic (m_1, \dots, m_t) -cycle system of $4K_v$, if and only if δ is a $(4K_v, \delta)$ -difference system.

The orbit of cycle C_n , denoted by $orb(C_n)$, is the set of all distinct n -cycles in the collection $\{C_n + i | i \in Z_v\}$. The length of $orb(C_n)$ is its cardinality, i.e., $orb(C_n) = k$, where k is the minimum positive integer such

that $C_n + k = C_n$. A cycle orbit of length ν on λK_ν is said to be *full* and otherwise *short*.

3. A Near Cyclic (m_1, m_2, \dots, m_r) -cycle System

In this section, we present new definitions and results of a near cyclic (m_1, m_2, \dots, m_r) -cycle system, that are useful for our proof.

Definition 3.1. A near cyclic (m_1, \dots, m_r) -cycle system of $4K_\nu$, $NCCS(4K_\nu, \delta)$, combining a near-four-factor and cyclic (m_1, \dots, m_r) -cycle system that is generated by the starter set δ . In addition, $NCCS(4K_\nu, \delta)$ is a $(\nu \times |\delta|)$ array that satisfies the following conditions:

- the cycles in row r and column i form a near-four-factor with focus r ,
- the cycles associated with rows contain no repetitions.

Undoubtedly, for presenting the $NCCS(4K_\nu, \delta)$, it is sufficient to provide a starter set δ that satisfied a near-four-factor.

We present here some of new definitions which will be needed in the sequel.

Definition 3.2. Two m -cycles H and F of a graph G of order ν are said to be *parallel* if they have the same difference set.

Definition 3.3. Let H and F be two m -cycles of a graph G of order ν . If the sum of each two corresponding vertices of them is ν , then it is called *adjoined m -cycles*, i.e., for $H = (h_1, h_2, \dots, h_m)$ and $F = (f_1, f_2, \dots, f_m)$ if $h_i + f_i = \nu, i = 1, \dots, m$, then H and F are adjoined cycles.

Corollary 3.1. Any two adjoined cycles are parallel cycles.

Throughout the paper, we shall sometimes use superscripts to identify the number of the cycles in a set. So, let us consider $\delta = \{C_{m_1}^{n_1}, C_{m_2}^{n_2}, \dots, C_{m_r}^{n_r}\}$ to be the set comprised of n_i cycles of length m_i , for $i = 1, 2, \dots, r$. In addition, we consider that C_{m_i} is the i th m -cycle in starter

set δ . Therefore, it is convenient to provide an example here to clarify the above discussion.

Example 3.1. Let $G = 4K_{22}$ and $\delta = \{C_4^5, C_{11}^2\}$ be a set of cycles of G such that

$$C_{4_1} = (1, 21, 12, 10), C_{4_2} = (2, 20, 13, 9), C_{4_3} = (3, 19, 14, 8),$$

$$C_{4_4} = (4, 18, 7, 15), C_{4_5} = (5, 17, 16, 6),$$

$$C_{11_1} = (2, 11, 3, 10, 4, 9, 6, 8, 7, 17, 21),$$

$$C_{11_2} = (20, 11, 19, 12, 18, 13, 16, 14, 15, 5, 1).$$

Firstly, we note that each nonzero element in Z_{22} occurs twice in the cycles of δ . So every vertex has a degree 4 except zero element (isolated vertex) has degree zero. So, it satisfies the near-four-factor. Secondly, the difference sets for the cycles in δ are listed in Table 3.1 and Table 3.2 for 4-cycles and 11-cycles, respectively.

Table 3.1

4-cycle	(1, 21, 12, 10)	(2, 20, 13, 9)	(3, 19, 14, 8)	(4, 18, 7, 15)	(5, 17, 16, 6)
Difference set	{2, 9, 2, 9}	{4, 7, 4, 7}	{6, 5, 6, 5}	{8, 11, 8, 11}	{10, 1, 10, 1}

Table 3.2

11-cycle	(2, 11, 3, 10, 4, 9, 6, 8, 7, 17, 21)	(20, 11, 19, 12, 18, 13, 16, 14, 15, 5, 1)
Difference set	{9, 8, 7, 6, 5, 3, 2, 1, 10, 4, 3}	{9, 8, 7, 6, 5, 3, 2, 1, 10, 4, 3}

As clearly shown, we observe that $D(\delta) = D\left(\bigcup_{i=1}^5 C_{4_i}\right) \cup D\left(\bigcup_{i=1}^2 C_{11_i}\right)$

covers each element of Z_{11}^* four times while the middle difference $\frac{22}{2} = 11$ appears exactly twice. Therefore, the set $\delta = \{C_4^5, C_{11}^2\}$ is a $(4K_{22}, \delta)$ -difference system. Then an $NCCS(4K_{22}, \delta)$ is (22×7) array and the starter set $\delta = \{C_4^5, C_{11}^2\}$ generates all the cycles in (22×7) array by repeated addition of 1 (mod 22) as shown in Table 3.3.

Table 3.3

Focus	NCCS($4K_v, \delta$)									
0	1 21 12 10	2 20 13 9	3 19 14 8	...	20 11 19 12 18 13 16 14 15 5 1					
1	2 0 13 11	3 21 14 10	4 20 15 9	...	21 12 20 13 19 14 17 15 16 6 2					
⋮	⋮	⋮	⋮	...	⋮					
20	21 19 10 8	0 18 11 7	1 17 12 6	...	18 9 17 10 16 11 14 12 13 3 21					
21	0 20 11 9	1 19 12 8	2 18 13 7	...	19 10 18 11 17 12 15 13 14 4 0					

As usual, any m -cycle has been written as a permutation

$$(a_{1,1}, \dots, a_{1,n}, a_{2,1}, \dots, a_{2,r}, a_{3,1}, \dots, a_{3,l}),$$

where $n + r + l = m$. For the sake of simplicity, it can be represented as connected paths, we mean that $C_m = (P_{1,n}, P_{2,r}, P_{3,l})$ such that $P_{1,n} = [a_{1,1}, \dots, a_{1,n}]$, $P_{2,r} = [a_{2,1}, \dots, a_{2,r}]$, $P_{3,l} = [a_{3,1}, \dots, a_{3,l}]$.

We will define the difference between any two paths H and K , denoted by $D(H, K)$, as the difference between the last vertex in the path H and the first vertex in the path K . Thus, for the cycle $C_m = (P_{1,n}, P_{2,r}, P_{3,l})$, we find that $D(P_{1,n}, P_{2,r}) = D([a_{1,n}, a_{2,1}])$, $D(P_{2,r}, P_{3,l}) = D([a_{2,r}, a_{3,1}])$ and $D(P_{3,l}, P_{1,n}) = D([a_{3,l}, a_{1,1}])$. Subsequently,

$$D(C_m) = D(P_{1,n}) \cup D(P_{2,r}) \cup D(P_{3,l}) \cup D(P_{1,n}, P_{2,r}) \\ \cup D(P_{2,r}, P_{3,l}) \cup D(P_{3,l}, P_{1,n})$$

and $V(C_m) = V(P_{1,n}) \cup V(P_{2,r}) \cup V(P_{3,l})$.

Now we are ready to present the proof for Theorem 1.1, the main aim of our paper. We distinguish two cases according to the congruence class of $v \equiv (\text{mod } 12)$.

Case 1. There exists a full near cyclic (m_1, \dots, m_r) -cycle system of $4K_{12n+10}$, $NCCS(4K_{12n+10}, \delta)$.

Proof. We have two subcases:

Subcase 1. n is odd.

Suppose $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is the starter set of $4K_{12n+10}$ such that the list of 4-cycles is:

$$\begin{aligned} C_{4_i} &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) \\ &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (i, 12n+10-i, 6n+5+i, 6n+5-i), \end{aligned}$$

when $i = \frac{5n+3}{2}$, let

$$C_{4_i} = \left(\frac{5n+3}{2}, 12n+10 - \frac{5n+3}{2}, 6n+5 - \frac{5n+3}{2}, 6n+5 + \frac{5n+3}{2} \right).$$

While we consider C_{6n+5}^* and C_{6n+5}^{**} that are adjoined $(6n+5)$ -cycle such that $C_{6n+5}^* = (P_1^*, P_2^*, P_3^*)$, $C_{6n+5}^{**} = (P_1^{**}, P_2^{**}, P_3^{**})$, where $\{P_i^*, P_i^{**} \mid 1 \leq i \leq 3\}$ are paths as follows:

$$P_1^* = [2, 6n+5, 3, 6n+4, \dots, 2n+2, 4n+5], P_2^* = [3n+3, 3n+5, 3n+4],$$

$$P_3^* = [9n+8, 9n+4, 9n+9, 9n+3, \dots, 8n+6, 10n+7, 12n+9],$$

$$P_1^{**} = [12n+8, 6n+5, 12n+7, 6n+6, \dots, 10n+8, 8n+5],$$

$$P_2^{**} = [9n+7, 9n+5, 9n+6],$$

$$P_3^{**} = [3n+2, 3n+6, 3n+1, 3n+7, \dots, 4n+4, 2n+3, 1].$$

We will divide the proof into two parts as follows:

Part 1. In this part, we prove that δ is a near-four-factor. To do this, we need to calculate the vertices

$$V\left(\bigcup_{i=1}^{3n+2} C_{4_i}\right) = c_{1,i} \cup c_{2,i} \cup c_{3,i} \cup c_{4,i}, 1 \leq i \leq 3n+2$$

such that $c_{1,i} = i$, $c_{2,i} = 12n + 10 - i$, $c_{3,i} = 6n + 5 + i$, $c_{4,i} = 6n + 5 - i$, $1 \leq i \leq 3n + 2$, $i \neq \frac{5n+3}{2}$. Then

$$c_{1,i} = \{1, 2, 3, \dots, 3n + 2\} - \left\{\frac{5n+3}{2}\right\},$$

$$c_{2,i} = \{12n + 9, 12n + 8, \dots, 9n + 8\} - \left\{\frac{19n+17}{2}\right\},$$

$$c_{3,i} = \{6n + 6, 6n + 7, \dots, 9n + 7\} - \left\{\frac{17n+13}{2}\right\},$$

$$c_{4,i} = \{6n + 4, 6n + 3, \dots, 3n + 3\} - \left\{\frac{7n+7}{2}\right\}.$$

While, if $i = \frac{5n+3}{2}$, then

$$V(C_{4_i}) = \left\{\frac{5n+3}{2}, \frac{19n+17}{2}, \frac{7n+7}{2}, \frac{17n+13}{2}\right\}.$$

Observe that the vertices of all 4-cycles cover every nonzero elements of $\{Z_{12n+10} - \{6n+5\}\}$ exactly once, whereas we provide the vertices of $(6n+5)$ -cycles as $V(P_i^*) \cup V(P_i^{**})$, $i = 1, 2, 3$ as follows:

$$V(P_1^*) = \{2, 3, 4, \dots, 2n + 2\} \cup \{6n + 5, 6n + 4, \dots, 4n + 5\},$$

$$V(P_2^*) = \{3n + 3, 3n + 5, 3n + 4\},$$

$$V(P_3^*) = \{9n + 8, 9n + 9, \dots, 10n + 7\}$$

$$\cup \{9n + 4, 9n + 3, \dots, 8n + 6\} \cup \{12n + 9\},$$

$$V(P_1^{**}) = \{12n + 8, 12n + 7, \dots, 10n + 8\} \cup \{6n + 5, 6n + 6, \dots, 8n + 5\},$$

$$V(P_2^{**}) = \{9n + 7, 9n + 5, 9n + 6\},$$

$$V(P_3^{**}) = \{3n + 2, 3n + 1, \dots, 2n + 3\} \cup \{3n + 6, 3n + 7, \dots, 4n + 4\} \cup \{1\}.$$

Then the vertices of $(6n + 5)$ -cycles cover each nonzero element of Z_{12n+10} exactly once except $\{6n + 5\}$ twice. Then the vertex set of the cycles in δ , $V(\delta)$, covers each element of Z_{12n+10}^* twice. Consequently, it satisfies near-four-factor (with isolated zero element).

Part 2. In this part, we prove that $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is the $(4K_{12n+10}, \delta)$ -difference system. So, we will check the difference as follows:

$$\bigcup_{i=1}^{3n+2} D(c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) = \bigcup_{i=1}^{3n+2} D(c_{j,i}, c_{j+1,i}), 1 \leq j \leq 4,$$

where $c_{5,i} = c_{1,i}$,

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} D(c_{1,i}, c_{2,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (2i) = \{2, 4, \dots, 6n + 4\} - \{5n + 3\},$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} D(c_{2,i}, c_{3,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (6n + 5 - 2i)$$

$$= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{n + 2\},$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} D(c_{3,i}, c_{4,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (2i) = \{2, 4, \dots, 6n + 4\} - \{5n + 3\},$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} D(c_{4,i}, c_{1,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+3}{2}}}^{3n+2} (6n + 5 - 2i)$$

$$= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{n + 2\}.$$

When $i = \frac{5n+3}{2}$, then $D(C_{4_i}) = \{5n+3, 6n+5, 5n+3, 6n+5\}$.

Then the list of difference set of 4-cycles covers every element of $\{Z_{6n+5}^* - (n+2)\} \cup \{6n+5\}$ exactly twice. Similarly, we compute $D(C_{6n+5}^*) \cup D(C_{6n+5}^{**})$ as follows:

$$D(C_{6n+5}^*) = D(P_1^*) \cup D(P_2^*) \cup D(P_3^*) \cup D(P_1^*, P_2^*) \cup D(P_2^*, P_3^*) \cup D(P_3^*, P_1^*),$$

$$D(P_1^*) = \{6n+3, 6n+2, \dots, 2n+4, 2n+3\}, D(P_2^*) = \{2, 1\},$$

$$D(P_3^*) = \{4, 5, \dots, 2n+1, 2n+2\},$$

$$D(P_1^*, P_2^*) = D(4n+5, 3n+3) = \{n+2\},$$

$$D(P_2^*, P_3^*) = D(3n+4, 9n+8) = \{6n+4\},$$

$$D(P_3^*, P_1^*) = D(12n+9, 2) = \{3\}.$$

Relying on adjoined cycles C_{6n+5}^{**} and C_{6n+5}^* , we find the same difference set by Corollary 3.1. Then $D(C_{6n+5}^*) \cup D(C_{6n+5}^{**})$ covers each element of Z_{6n+5}^* exactly twice except $\{n+2\}$ four times. From the above discussion, we deduce that $D(\delta)$ covers each element in Z_{6n+5}^* four times and the middle difference $\{6n+5\}$ twice.

This assures that $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is $(4K_{12n+10}, \delta)$ -difference system, n is odd. Therefore, $\delta = \{C_4^{3n+2}, C_{6n+1}^2\}$ is starter set for the $NCCS(4K_{12v+10}, \delta)$ when n is odd. \square

Subcase 2. n is even.

Suppose $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is the starter set of $4K_{12n+10}$ such that the list of 4-cycles is:

$$\begin{aligned}
C_{4_i} &= \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) \\
&= \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (i, 12n + 10 - i, 6n + 5 + i, 6n + 5 - i).
\end{aligned}$$

When $i = \frac{n}{2}$, then $C_{4_i} = \left(\frac{n}{2}, 6n + 5 - \frac{n}{2}, 12n + 10 - \frac{n}{2}, 6n + 5 + \frac{n}{2}\right)$

whereas C_{6n+5}^* and C_{6n+5}^{**} are adjoined $(6n + 5)$ -cycles such that $C_{6n+5}^* = (P_1^*, P_2^*, P_3^*)$, $C_{6n+5}^{**} = (P_1^{**}, P_2^{**}, P_3^{**})$, where $\{P_i^*, P_i^{**} \mid 1 \leq i \leq 3\}$ are paths as follows:

$$P_1^* = [2, 6n + 5, 3, 6n + 4, \dots, 2n + 2, 4n + 5],$$

$$P_2^* = [3n + 5, 3n + 3, 3n + 4],$$

$$P_3^* = [9n + 8, 9n + 4, 9n + 9, 9n + 3, \dots, 8n + 6, 10n + 7, 12n + 9],$$

$$P_1^{**} = [12n + 8, 6n + 5, 12n + 7, 6n + 6, \dots, 10n + 8, 8n + 5],$$

$$P_2^{**} = [9n + 5, 9n + 7, 9n + 6],$$

$$P_3^{**} = [3n + 2, 3n + 6, 3n + 1, 3n + 7, \dots, 4n + 4, 2n + 3, 1].$$

In similar way for the Subcase 1, one may easily verify that $V(\delta) = \left(V\left(\bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} C_{4_i}\right) \cup V(C_{6n+5}^*) \cup V(C_{6n+5}^{**})\right)$ covers each element in Z_{12n+10}^* exactly twice. Now, we are going to calculate the difference set of 4-cycles as follows:

$$\bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{j,i}, c_{j+1,i}), 1 \leq j \leq 4,$$

where $c_{5,i} = c_{1,i}$,

$$\bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{1,i}, c_{2,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (2i) = \{2, 4, \dots, 6n + 4\} - \{n\},$$

$$\begin{aligned} \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{2,i}, c_{3,i}) &= \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (6n + 5 - 2i) \\ &= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{5n + 5\}, \end{aligned}$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{3,i}, c_{4,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (2i) = \{2, 4, \dots, 6n + 4\} - \{n\},$$

$$\begin{aligned} \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} D(c_{4,i}, c_{1,i}) &= \bigcup_{\substack{i=1 \\ i \neq \frac{n}{2}}}^{3n+2} (2i6n + 5 - 2i) \\ &= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{5n + 5\}. \end{aligned}$$

When $i = \frac{n}{2}$, $D(C_{4_i}) = \{5n + 5, 6n + 5, 5n + 5, 6n + 5\}$.

Then the list of difference set of 4-cycles covers each element of $\{Z_{6n+5}^* - (n)\} \cup \{6n + 5\}$ exactly twice. Correspondingly, the list of difference set of $(6n + 5)$ -cycles calculates as follows:

$$\begin{aligned} D(C_{6n+5}^*) &= D(P_1^*) \cup D(P_2^*) \cup D(P_3^*) \cup D(P_1^*, P_2^*) \\ &\quad \cup D(P_2^*, P_3^*) \cup D(P_3^*, P_1^*), \end{aligned}$$

$$D(P_1^*) = \{6n + 3, 6n + 2, \dots, 2n + 4, 2n + 3\}, D(P_2^*) = \{2, 1\},$$

$$D(P_3^*) = \{4, 5, \dots, 2n + 1, 2n + 2\}, D(P_1^*, P_2^*) = D(4n + 5, 3n + 5) = \{n\},$$

$$D(P_2^*, P_3^*) = D(3n + 4, 9n + 8) = \{6n + 4\}, D(P_3^*, P_1^*) = D(12n + 9, 2) = \{3\}.$$

As clearly shown, in the previous equation, the vertices of $6n + 5$ -cycles cover every element of Z_{6n+5}^* exactly twice except $\{n\}$ four times. Thus,

we realize now that $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is $(4K_{12n+10}, \delta)$ -difference system, n is even. Then $\delta = \{C_4^{3n+2}, C_{6n+5}^2\}$ is starter set for the $NCCS(4K_{12v+10}, \delta)$ when n is even. \square

Case 2. There exists a full cyclic (m_1, \dots, m_r) -cycle system of $4K_{12n+2}, NCCS(4K_{12n+2}, \delta)$.

Proof. We also have two subcases:

Subcase 1. n is even.

When $n = 2$, $v = 26$, let $\delta = \{C_4^6, C_7^2, C_6^2\}$ be the starter set of $NCCS(4K_{26}, \delta)$ as follows:

$$C_{4_1} = (1, 25, 14, 12), C_{4_2} = (2, 24, 15, 11), C_{4_3} = (3, 23, 16, 10),$$

$$C_{4_4} = (4, 22, 17, 9), C_{4_5} = (5, 21, 18, 8), C_{4_6} = (6, 19, 7, 20),$$

$$C_7^* = (13, 2, 12, 3, 11, 4, 10), C_7^{**} = (13, 24, 14, 23, 15, 22, 16),$$

$$C_6^* = (6, 1, 5, 17, 19, 18), C_6^{**} = (20, 25, 21, 9, 7, 8).$$

It is straightforward to check that δ is actually a starter set of $NCCS(4K_{26}, \delta)$.

When $n \geq 4$, suppose $\delta = \{C_4^{3n}, C_{2n+2}^2, C_{4n-1}^2\}$ is the starter set of $NCCS(4K_{12n+2}, \delta)$ such that the list of 4-cycles is:

$$\begin{aligned} C_{4_i} &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) \\ &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (i, 12n+2-i, 6n+1+i, 6n+1-i), \end{aligned}$$

when $i = \frac{5n+4}{2}$ let

$$C_{4_i} = \left(\frac{5n+4}{2}, 6n+1 - \frac{5n+4}{2}, 12n+2 - \frac{5n+4}{2}, 6n+1 + \frac{5n+4}{2} \right).$$

While we consider C_{4n-1}^* and C_{4n-1}^{**} that are adjoined $(4n-1)$ -cycles such that

$$C_{4n-1}^* = (6n+1, 2, 6n, 3, 6n-1, 4, \dots, 2n-1, 4n+3, 2n, 4n+2),$$

$$C_{4n-1}^{**} = (6n+1, 12n, 6n+2, 12n-1, 6n+3, \dots, 10n+3, 8n-1, 10n+2, 8n).$$

As well, we consider that C_{2n+2}^* and C_{2n+2}^{**} are adjoined $(2n+2)$ -cycles such that

$$\begin{aligned} &C_{2n+2}^* \\ &= (2n+2, 1, 2n+1, 8n+1, 10n-1, 8n+2, 10n-2, \dots, 9n+2, 9n-1, 9n+1, 9n), \end{aligned}$$

$$\begin{aligned} &C_{2n+2}^{**} \\ &= (10n, 12n+1, 10n+1, 4n+1, 2n+3, 4n, 2n+4, \dots, 3n, 3n+3, 3n+1, 3n+2). \end{aligned}$$

Similarly, it will be following the same manner of the previous case to prove that the set δ is the starter set of $4K_{12n+2}$. We will divide the proof into two parts as follows:

Part 1. In this part, we prove a near-four-factor. So, we need to calculate the vertices $V\left(\bigcup_{i=1}^{3n} C_{4_i}\right) = c_{1,i} \cup c_{2,i} \cup c_{3,i} \cup c_{4,i}, 1 \leq i \leq 3n$ such that

$$c_{1,i} = i, c_{2,i} = 12n+2-i, c_{3,i} = 6n+1+i,$$

$$c_{4,i} = 6n+1-i, 1 \leq i \leq 3n+2, i \neq \frac{5n+4}{2}.$$

$$c_{1,i} = \{1, 2, 3, \dots, 3n\} - \left\{ \frac{5n+4}{2} \right\}, c_{2,i} = \{12n+1, 12n, \dots, 9n+2\} - \left\{ \frac{19n}{2} \right\},$$

$$c_{3,i} = \{6n+2, 6n+3, \dots, 9n+1\} - \left\{ \frac{17n+6}{2} \right\},$$

$$c_{4,i} = \{6n, 6n - 1, \dots, 3n + 1\} - \left\{ \frac{7n - 2}{2} \right\}.$$

$$\text{And when } i = \frac{5n + 4}{2}, \text{ then } V(C_{4_i}) = \left\{ \frac{5n + 4}{2}, \frac{7n - 2}{2}, \frac{19n}{2}, \frac{17n + 6}{2} \right\}.$$

At the same time, the vertex set of remaining cycles can be written as follows:

$$V(C_{4n-1}^*) = \{2, 3, 4, \dots, 2n\} \cup \{4n + 2, 4n + 3, \dots, 6n + 1\},$$

$$V(C_{4n-1}^{**}) = \{6n + 1, 6n + 2, \dots, 8n\} \cup \{10n + 2, 10n + 3, \dots, 12n\},$$

$$V(C_{2n+2}^*) = \{1, 2n + 1, 2n + 2\} \cup \{8n + 1, 8n + 2, 8n + 3, \dots, 10n - 2, 10n - 1\},$$

$$V(C_{2n+2}^{**}) = \{12n + 1, 10n, 10n + 1\} \cup \{2n + 3, 2n + 4, 2n + 5, \dots, 4n, 4n + 1\}.$$

Simply we can note that $V(\delta)$ covers $\{Z_{12n+2}^*\}$ exactly twice.

Part 2. In this part, we prove that $\delta = \{C_4^{3n}, C_{4n-1}^2, C_{2n+2}^2\}$ is the $(4K_{12n+2}, \delta)$ -difference system. So, we check the difference as follows:

The list of difference set of all 4-cycles $\left(\bigcup_{i=1}^{3n} D(C_{4_i}) \right)$ is determined as follows:

$$\bigcup_{i=1}^{3n} D(C_{4_i}) = \bigcup_{i=1}^{3n} D(c_{j,i}, c_{j+1,i}), 1 \leq j \leq 4, \text{ where } c_{5,i} = c_{1,i},$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} D(c_{1,i}, c_{2,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (2i) = \{2, 4, \dots, 6n\} - \{5n + 4\},$$

$$\begin{aligned} \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} D(c_{2,i}, c_{3,i}) &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (6n + 1 - 2i) \\ &= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{n - 3\}, \end{aligned}$$

$$\bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} D(c_{3,i}, c_{4,i}) = \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (2i) = \{2, 4, \dots, 6n\} - \{5n + 4\},$$

$$\begin{aligned} \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} D(c_{4,i}, c_{1,i}) &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+4}{2}}}^{3n} (6n + 1 - 2i) \\ &= \{6n + 3, 6n + 1, \dots, 3, 1\} - \{n - 3\}. \end{aligned}$$

Also, when $i = \frac{5n + 4}{2}$, $D(C_{4_i}) = \{n - 3, 6n + 1, n - 3, 6n + 1\}$.

Then the list of difference set of all 4-cycles $(D(C_4^{3n}))$ covers each element of $\{Z_{6n+1}^* - (5n + 4)\} \cup \{6n + 1\}$ precisely twice. Correspondingly, the list of difference set of remaining cycles $\{C_{2n+2}^*, C_{2n+2}^{**}, C_{4n-1}^*, C_{4n-1}^{**}\}$ is computed as below:

$$D(C_{4n-1}^*) = D\{(6n + 1, 2, 6n, 3, 6n - 1, 4, \dots, 2n - 1, 4n + 3, 2n, 4n + 2)\},$$

$$D(C_{4n-1}^{**}) = \{6n - 1, 6n - 2, 6n - 3, \dots, 2n + 3, 2n + 2\} \cup \{2n - 1\}.$$

Since C_{4n-1}^* and C_{4n-1}^{**} are adjoined cycles in $4K_{12n+2}$, $D(C_{4n-1}^{**}) = D(C_{4n-1}^*)$.

We also have:

$$\begin{aligned} D(C_{2n+2}^*) &= D\{(2n + 2, 1, 2n + 1, 8n + 1, 10n - 1, 8n + 2, \\ &\quad 10n - 2, \dots, 9n + 2, 9n - 1, 9n + 1, 9n)\} \\ &= \{2n + 1, 2n, 6n, 2n - 2, 2n - 3, 2n - 4, \dots, 3, 2, 1\} \cup \{5n + 4\}. \end{aligned}$$

Since C_{2n+2}^* and C_{2n+2}^{**} are adjoined cycles in $4K_{12n+2}$, $D(C_{2n+2}^{**}) = D(C_{2n+2}^*)$.

Thus, each element in the multiset Z_{6n+1}^* is covered by $D(C_{4n-1}^*) \cup D(C_{4n-1}^{**}) \cup D(C_{2n+2}^*) \cup D(C_{2n+2}^{**})$ twice except $\{5n+4\}$ four times. In view of previous observation, we conclude that $\delta = \{C_4^{3n}, C_{2n+2}^2, C_{4n-1}^2\}$ is $(4K_{12n+2}, \delta)$ -difference system, n is even. \square

Subcase 2. n is odd.

Suppose $\delta = \{C_4^{3n}, C_{2n+2}^2, C_{4n-1}^2\}$ is the starter set of cycles of $NCCS(4K_{12n+2}, \delta)$ such that the list of 4-cycles is:

$$\begin{aligned} C_{4_i} &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+1}{2}}}^{3n} (c_{1,i}, c_{2,i}, c_{3,i}, c_{4,i}) \\ &= \bigcup_{\substack{i=1 \\ i \neq \frac{5n+1}{2}}}^{3n} (i, 12n+2-i, 6n+1+i, 6n+1-i), \end{aligned}$$

when $i = \frac{5n+1}{2}$, let

$$C_{4_i} = \left(\frac{5n+1}{2}, 12n+2 - \frac{5n+1}{2}, 6n+1 - \frac{5n+1}{2}, 6n+1 + \frac{5n+1}{2} \right)$$

whereas that C_{4n-1}^* and C_{4n-1}^{**} are adjoined $(4n-1)$ -cycles such that

$$C_{4n-1}^* = (6n+1, 2, 6n, 3, 6n-1, 4, \dots, 2n-1, 4n+3, 2n, 4n+2),$$

$$C_{4n-1}^{**} = (6n+1, 12n, 6n+2, 12n-1, 6n+3, \dots, 10n+3, 8n-1, 10n+2, 8n).$$

Also, we consider that C_{2n+2}^* and C_{2n+2}^{**} are adjoined $(2n+2)$ -cycles such that $C_{2n+2}^* = (P_1^*, P_2^*)$, $C_{2n+2}^{**} = (P_1^{**}, P_2^{**})$, where $\{P_i^*, P_i^{**} \mid 1 \leq i \leq 2\}$ are paths as follows:

$$P_1^* = [2n+2, 1, 10n+1],$$

$$P_2^* = [4n + 1, 2n + 3, 4n, 2n + 4, \dots, 3n, 3n + 3, 3n + 1, 3n + 2],$$

$$P_1^{**} = [10n, 12n + 1, 2n + 1],$$

$$P_2^{**} = [8n + 1, 10n - 1, 8n + 2, 10n - 2, \dots, 9n + 2, 9n - 1, 9n + 1, 9n].$$

Obviously, as the Subcase 1, it can be found that $V(\delta)$ covers each element of Z_{12n+2}^* exactly twice and the list of difference set of all 4-cycles $(D(C_4^{3n}))$ covers each element of $\{Z_{6n+1}^* - n\}$ precisely twice, whereas the difference set of $(4n - 1)$ -cycles $(D(C_{4n-1}^*) \cup D(C_{4n-1}^{**}))$ contains elements $\{6n - 1, 6n - 2, 6n - 3, \dots, 2n + 3, 2n + 2\} \cup \{2n - 1\}$ twice. Now, we calculate the difference set of $(2n + 2)$ -cycles as follows:

$$D(C_{2n+2}^*) = D(P_1^*) \cup D(P_2^*) \cup D(P_1^*, P_2^*) \cup D(P_2^*, P_1^*),$$

$$D(P_1^*) = \{2n + 1, 2n\}, D(P_2^*) = \{2n - 2, 2n - 3, 2n - 4, \dots, 3, 2, 1\},$$

$$D(P_1^*, P_2^*) = D(10n + 1, 4n + 1) = \{6n\}, D(P_2^*, P_1^*) = D(2n + 2, 3n + 2) = \{n\}.$$

Then all elements in the set $\{1, 2, 3, \dots, 2n - 3, 2n - 2, 2n, 2n + 1, 6n\}$ appear in $D(C_{2n+2}^*)$ exactly once except $\{n\}$ twice. Therefore, the multiset of $D(C_{4n-1}^*) \cup D(C_{4n-1}^{**}) \cup D(C_{2n+2}^*) \cup D(C_{2n+2}^{**})$ covers each element of $\{Z_{6n+1}^*\}$ exactly twice except $\{n\}$ four times.

Hence, $\delta = \{C_4^{3n}, C_{2n+2}^2, C_{4n-1}^2\}$ is $(4K_{12n+2}, \delta)$ -difference system, n is odd. Then $\delta = \{C_4^{3n}, C_{2n+2}^2, C_{4n-1}^2\}$ is starter set of $NCCS(4K_{12n+2}, \delta)$.

□

Acknowledgement

The authors thank the anonymous referees for their numerous suggestions and comments which improved the original manuscript.

References

- [1] B. Alspach, Research problems, *Discrete Math.* 36(3) (1981), 333-334.
- [2] B. Alspach and H. Gavlas, Cycle decompositions of K_n and $K_n - I$, *J. Combin. Theory Ser. B* 81(1) (2001), 77-99.
- [3] D. Bryant, D. Horsley and W. Pettersson, Cycle decompositions V: Complete graphs into cycles of arbitrary lengths, *Proc. London Math. Soc.*, 2013, doi: 10.1112/plms/pdt051.
- [4] D. Bryant, D. Horsley, B. Maenhaut and B. R. Smith, Decompositions of complete multigraphs into cycles of varying lengths, 4 August 2015, arXiv: 1508.00645v1 [math.CO].
- [5] M. Buratti, A description of any regular or 1-rotational design by difference methods, *Booklet of the Abstracts of Combinatorics*, 2000, pp. 35-52.
- [6] A. Kotzig, Decompositions of a complete graph into $4k$ -gons, *Matematický Časopis* 15 (1965), 229-233.
- [7] K. Matarneh and H. Ibrahim, Array cyclic $(5^*, 6^{**}, 4)$ -cycle design, *Far East J. Math. Sci. (FJMS)* 100(10) (2016), 1611-1626.
- [8] R. Pelsesohn, Eine Lösung der beiden Heffterschen Differenzenprobleme, *Compos. Math.* 6 (1939), 251-257.
- [9] A. Rosa, On cyclic decompositions of the complete graph into $(4m + 2)$ -gons, *Matematicko-Fyzikálny Časopis* 16(4) (1966), 349-352.
- [10] A. Rosa, On the cyclic decompositions of the complete graph into polygons with an odd number of edges, *Časopis Pest. Math.* 91 (1966), 53-63.
- [11] M. Šajna, Cycle decompositions III: complete graphs and fixed length cycles, *J. Combin. Des.* 10(1) (2002), 27-78.